

ISSN 0017 - 8861

Estd 1923

Health

A JOURNAL DEVOTED TO HEALTHFUL LIVING

Email: admin@theantiseptic.in
subscription@theantiseptic.in

www.healthjournal.in

Vol. 104 □ No. 07

JULY 2026

Rs. 20

Hoarseness



Page No : 32

TINNITUS AND ROLE OF ANTIOXIDANTS

PART - 2

(Continued from previous issue)

Antioxidants: The Why

Increased oxidative stress and chronic inflammation are involved in hearing disorders. Since antioxidants reduce oxidative stress by neutralizing free radicals and reducing inflammation, supplementation with antioxidants may improve current prevention programs that use physical devices and improve the efficacy of standard therapy in patients with hearing disorders. Several studies with antioxidants and hearing disorders discussed above support this rationale.

Although supplementation with a single antioxidant in these studies has produced some beneficial effects in improving hearing disorders, this is not recommended for optimal results. This is due to the fact that an individual antioxidant, when oxidized, acts as a free radical. In addition, the use of a single antioxidant, such as vitamin E or beta-carotene in humans with a high internal oxidative environment (eg, heavy tobacco smokers), has produced harmful effects. This is consistent with the fact that individual antioxidants in high oxidative environments acted as a pro-oxidant. It is interesting to point

out that, unlike studies on high-risk human populations, animal studies with individual antioxidants consistently produced beneficial effects.

Antioxidants: The When and How

In addition to physical protection, antioxidants should be consumed daily prior to exposure to agents or conditions that induce hearing disorders to reduce the risk of developing tinnitus, hearing loss, or a balancing problem. People suffering from hearing disorders should take antioxidants in combination with standard therapy.

However, the selection of the appropriate type of antioxidant preparation and dose-schedule is critical for enhancing their effectiveness. The authors recommend supplementation with multiple micronutrients containing dietary and endogenous antioxidants for reducing the risk of hearing disorders and improving the efficacy of standard therapy to improve the management of hearing disorders.

A closer look at specific antioxidants.

Almost all antioxidants can act as pro-oxidants (free radicals) when oxidized. Therefore, as

mentioned earlier, the use of single antioxidants is not recommended. Multiple antioxidants are necessary because their organ and cellular distributions, as well as mechanisms of actions, are different.

For example, beta-carotene (BC) is more effective in quenching oxygen radicals than most other antioxidants. BC can perform certain biological functions that cannot be produced by its metabolite vitamin A, and vice versa. It has been reported that BC treatment enhances the expression of the connexin gene, which codes for a gap junction protein in mammalian fibroblasts in culture, whereas vitamin A treatment does not produce such an effect. Vitamin A induces differentiation in certain normal cells and cancer cells, whereas BC and other carotenoids do not. Thus, BC and vitamin A have, in part, different biological functions. The gradient of oxygen pressure varies within cells. Some antioxidants, such as vitamin E, are more effective as quenchers of free radicals in reduced oxygen pressure, whereas others, such as BC and vitamin A, are more effective in higher atmospheric pressures. Vitamin C is necessary to protect cellular components in aqueous environments, whereas carotenoids and vitamins A and E protect cellular components in lipid environments. In addition, vitamin C is necessary for the activity of tyrosine hydroxylase, which is the rate-limiting enzyme in the synthesis of catecholamines.



Oxidized forms of vitamin C and vitamin E can also act as radicals; therefore, excessive amounts of any one of these forms, when used as a single agent, could be harmful over a long period of time. Vitamin C also plays an important role in maintaining cellular levels of vitamin E by recycling vitamin E radical (oxidized) to the reduced (antioxidant) form. Also, oxidative DNA damage produced by levels of vitamin C could be protected by vitamin E.

The form of vitamin E used is also important in any clinical trial. It has been established that d-alpha-tocopheryl succinate (a-TS) is the most effective form of vitamin E both in vitro and in vivo. This form of vitamin E is more soluble than a-tocopherol and enters cells more readily. Therefore, it is expected to cross the blood-brain barrier in greater amounts than a-tocopherol. However, this has not yet to be demonstrated in animals or humans.

We have reported that oral ingestion of aTS (800 IU/day) in humans increased plasma levels of not only a-tocopherol, but also a-TS, suggesting that

a-TS can be absorbed from the intestinal tract before hydrolysis to a-tocopherol. This observation is important because the conventional assumption, based on rodents, has been that esterified forms of vitamin E such as a-tocopheryl acetate, a-tocopheryl nicotinate, and a-TS can be absorbed from the intestinal tract only after they are hydrolyzed to a-tocopherol. Our data show that this assumption may not be true for the absorption of a-TS in humans.

Another antioxidant is glutathione, which is effective in catabolizing H₂O₂ and anions. However, oral supplementation with glutathione failed to significantly increase plasma levels of glutathione in human subjects, suggesting that this tripeptide is completely hydrolyzed in the gastrointestinal (GI) tract. N-acetylcysteine and alpha-lipoic acid (glutathione-elevating agents by different mechanisms) can be used as an antioxidant in combination with others.

Besides well-characterized antioxidants—such as vitamin A, carotenoids, vitamin C, and vitamin E—other antioxidants are also important in protecting against cellular damage in the ear, including coenzyme and glutathione. A study has shown that ubiquinol (coenzyme Q10) scavenges peroxy radicals faster than a-tocopherol and, like vitamin C, can regenerate vitamin E in a redox cycle. However, it is a weaker antioxidant than

a-tocopherol. Selenium is a co-factor of glutathione peroxidase, and S-glutathione peroxidase also acts as an antioxidant. Therefore, selenium supplementation together with other antioxidants is also essential. In addition to antioxidants, B-vitamins, vitamin D, and certain minerals are essential for optimal health. The pertinent references for this section are described in recent reviews.^{44,45}

An antioxidant formulation containing iron or copper is not recommended, because these components interact with vitamin C and generate excessive amounts of free radicals. In addition, these trace minerals in the presence of vitamin C are absorbed better than in its absence, and nature has provided no significant mechanisms of excretion of iron or copper in men of all ages and women after menopause. Increased free iron or copper stores in the body increase the risk of many chronic diseases. The addition of heavy metals such as molybdenum, zirconium, and vanadium is also not recommended, because there are no significant mechanisms of removal of these heavy metals from the body. An accumulation of these metals after long-term consumption could be toxic to nervous tissue including the brain.

The inclusion of herbs or herbal antioxidants into a multiple dietary and endogenous antioxidant preparation is not recommended,

because they do not produce any unique beneficial effects that cannot be produced by standard antioxidants. In addition, certain herbs are known to interact with prescription and non-prescription drugs in an adverse manner. These issues have been discussed in detail in two review papers.^{44,45}

How much? Doses are very important because, at certain low doses, antioxidants may reduce free radicals but may not decrease chronic inflammation. At higher doses (totally safe in humans), they reduce oxidative stress as well as inflammation. These issues have been discussed in detail in a recent review paper.⁴⁴

A dose-schedule of twice a day (half in the morning and half in the evening) is equally important for increasing effectiveness. Taking antioxidants once a day may not be adequate, because of the resultant fluctuations of antioxidant levels in the body. For example, if antioxidants are consumed in the morning, at least half have been eliminated from the body by evening, and at least another half by the next morning. This can create significant fluctuations in antioxidant levels; only a twofold change in the level of vitamin E succinate is known to cause marked alterations in gene expression. Thus, the genetic machinery of the cell constantly has to readjust to cope with this fluctuation in the levels of antioxidants, and this

could create cellular stress over a long period of time.⁴⁵

Clinical Trial of Antioxidant Formulation for Hearing Health

Using the concepts of antioxidant formulation described above, PMC has prepared a formulation referred to as Hearing Health for preventing and improving hearing disorders in combination with standard therapy.

PMC antioxidant formulation was tested in a clinical study in troops returning from Iraq with mild to moderate traumatic brain injury. A total of 34 patients with post-traumatic dizziness were admitted to the Naval Medical Center San Diego Clinic over a 2-month period and the patients agreed to participate in the study under the supervision of Dr Michael Hoffer and his colleagues.

All patients had received their injury 3 to 20 weeks prior to admission, and they received identical treatment consisting of medical therapy (for any migraines), supportive care, steroids, and vestibular rehabilitation therapy. A total of 15 of the 34 patients also received a dose of an antioxidant and micronutrient formula (two capsules by mouth twice a day). At the onset of therapy, all patients were evaluated in outcome measures, which included the Sensory Organization Test (SOT) by Computerized Dynamic Posturography (CDP), the Dynamic

Gait Index (DGI), the Activities Balance Confidence (ABC) scale, the Dizziness Handicap Index (DHI), the Vestibular Disorders Activities of Daily Living (VADL) score, and the Balance Scoring System (BESS) test.

The study was carried out for 12 weeks. The therapist who graded these outcomes and performed the testing was blinded as to whether the patient was receiving antioxidant therapy or not. The pretrial test scores did not differ significantly between the two groups on any of the tests.

Both groups of patients showed trends toward significant improvement on all tests after the 12 weeks of therapy, but the combination treatment trend was stronger than that of the standard therapy alone group. After only 4 weeks, the SOT score by CDP was 78 for the antioxidant group as compared to 63 for the non-antioxidant group. This difference was statistically significant at the $P < 0.05$ level. The improvement noted by the antioxidant group on the other tests was also greater than the non-antioxidant group, although these differences did not reach statistical significance because of the short trial period and small sample size.


In addition to the above clinical trial, several consumers who are on PMC Hearing Health formulation are showing improvement in hearing disorders. Almost all previous

studies using primarily one, and occasionally two, components showed some beneficial effects in improving hearing disorders, but the PMC Hearing Health formula is very comprehensive and includes (at the proper dose, type, and dose schedule) dietary and endogenous antioxidants, B-vitamins, vitamin D, selenomethionine, and other minerals, but no iron, copper, or manganese or heavy metals or herbal products. This product is available commercially (www.mypmcinside.com).

References

1. Diao MF, Liu HY, Zhang YM, Gao WY [Changes in antioxidant capacity of the guinea pig exposed to noise and the protective effect of alpha-lipoic acid against acoustic trauma]. *Sheng Li Xue Bao.* 2003;55(6):672-6.
2. Clerici WJ, DiMartino DL, Prasad MR. Direct effects of reactive oxygen species on cochlear outer hair cell shape in vitro. *Hear Res.* 1995;84(1-2):30-40.
3. Henderson D, Bielefeld EC, Harris KC, Hu BH. The role of oxidative stress in noise-induced hearing loss. *Ear Hear.* 2006;27(1):1-19.
4. Henderson D, McFadden SL, Liu CC, Hight N, Zheng X Y. The role of antioxidants in protection from impulse noise. *Ann NY Acad Sci.* 1999;884:368-80.
5. Ohlemiller KK, Wright JS, Dugan LL. Early elevation of cochlear reactive oxygen species following noise exposure. *Audiol Neurootol.* 1999;4(5):229-36.

6. Van Campen LE, Murphy WJ, Franks JR, Mathias PI, Toraason MA. Oxidative DNA damage is associated with intense noise exposure in the rat. *Hear Res.*2002;164(1-2):29-38.
7. Yamane H, Nakai Y, Takayama M, et al. The emergence of free radicals after acoustic trauma and strial blood flow. *Acta Otolaryngol Suppl.*1995;519:87-92.
8. Yamashita D, Jiang H Y, Schacht J, Miller JM. Delayed production of free radicals following noise exposure. *Brain Res.*2004;1019(1-2):201-9.
9. Zou J, Bretlau P, Pyykko I, Starck J, Toppila E. Sensorineural hearing loss after vibration: an animal model for evaluating prevention and treatment of inner ear hearing loss. *Acta Otolaryngol.*2001;121(2):143-8.
10. Minami SB, Sha SH, Schacht J. Antioxidant protection in a new animal model of cis-platin-induced ototoxicity. *Hear Res.* 2004;198(1-2):137-43.
11. Husain K, Scott RB, Whitworth C, Somani SM, Rybak L P. Dose response of carboplatin-induced hearing loss in rats: antioxidant defense system. *Hear Res.*2001;151(1-2):71-78.
12. Neri S, Signorelli S, Pulvirenti D, et al. Oxidative stress, nitric oxide, endothelial dysfunction and tinnitus. *Free Radic Res.* 2006;40(6):615-8.
13. Pall ML, Bedient SA. The NO/ONOO-cycle as the etiological mechanism of tinnitus. *Int Tinnitus J.*2007;13(2):99-104.
14. Kowalska S, Sulkowski W. [Tinnitus in noise-induced hearing impairment]. *Med Pr.* 2001;52(5):305-13.
15. Fagelson MA. The association between tinnitus and posttraumatic stress disorder. *Am J Audiol.*2007;16(2):107-17.
16. Takumida M, Anniko M, Ohtani M. Radical scavengers for Meniere's disease after failure of conventional therapy: a pilot study. *Acta Otolaryngol.*2003;123(6):697-703.
17. Le T, Keithley EM. Effects of antioxidants on the aging inner ear. *Hear Res.* 2007;226(1-2):194-202.
18. Hou F, Wang S, Zhai S, Hu Y, Yang W, He L. Effects of alpha-tocopherol on noise-induced hearing loss in guinea pigs. *Hear Res.* 2003;179(1-2):1-8.
19. Joachims HZ, Segal J, Golz A, Netzer A, Goldenberg D. Antioxidants in treatment of idiopathic sudden hearing loss. *Otol Neurotol.*2003;24(4):572-5.
20. Gordin A, Goldenberg D, Golz A, Netzer A, Joachims HZ. Magnesium: a new therapy for idiopathic sudden sensorineural hearing loss. *Otol Neurotol.*2002;23(4):447-51.
21. Angeli SI, Liu XZ, Yan D, Balkany T, Telischi F. Coenzyme Q-10 treatment of patients with a 7445A>G mitochondrial DNA mutation stops the progression of hearing loss. *Acta Otolaryngol.*2005;125(5):510-2.
22. Seidman MD. Glutamate antagonists, steroids, and antioxidants as therapeutic options for hearing loss and tinnitus and the use of an inner ear drug delivery system. *Int Tinnitus J.*1998;4(2):148-154.
23. Scholik AR, Lee US, Chow CK, Yan H Y. Dietary vitamin E protects the fathead minnow, *Pimephales promelas*, against noise exposure. *Comp Biochem Physiol C Toxicol Pharmacol.*2004;137(4):313-23.
24. Duan M, Qiu J, Laurell G, Olofsson A, Counter SA, Borg E. Dose and time-dependent protection of the antioxidant N-L-acetyl-cysteine against impulse noise trauma. *Hear Res.*2004;192(1-2):1-9.
25. Ohinata Y, Miller JM, Schacht J. Protection from noise-induced lipid peroxidation and hair cell loss in the cochlea. *Brain Res.*2003;966(2):265-73.
26. Kopke R, Bielefeld E, Liu J, et al. Prevention of impulse noise-induced hearing loss with antioxidants. *Acta Otolaryngol.* 2005;125(3): 235-43.

27. Sato K. Pharmacokinetics of coenzyme Q10 in recovery of acute sensorineural hearing loss due to hypoxia. *Acta Otolaryngol Suppl.* 1988;458:95-102.
28. Sergi B, Fetoni AR, Paludetti G, et al. Protective properties of idebenone in noise-induced hearing loss in the guinea pig. *Neuroreport.*2006;17(9):857-61.
29. McFadden SL, Woo JM, Michalak N, Ding D. Dietary vitamin C supplementation reduces noise-induced hearing loss in guinea pigs. *Hear Res.*2005;202(1-2):200-8.
30. Kalkanis JG, Whitworth C, Rybak L P. Vitamin E reduces cisplatin ototoxicity. *Laryngoscope.* 2004;114(3):538-42.
31. Teranishi MA, Nakashima T. Effects of trolox, locally applied on round windows, on cis-platin-induced ototoxicity in guinea pigs. *Int J Pediatr Otorhinolaryngol.*2003;67(2):133-9.
32. Sergi B, Fetoni AR, Ferraresi A, et al. The role of antioxidants in protection from ototoxic drugs. *Acta Otolaryngol Suppl.*2004 May;(552):42-5.
33. Fetoni AR, Sergi B, Ferraresi A, Paludetti G, Troiani D. alpha-Tocopherol protective effects on gentamicin ototoxicity: an experimental study. *Int J Audiol.*2004;43(3):166-71.
34. Feghali JG, Liu W, Van De Water TR. L-n-acetyl-cysteine protection against cisplatin-induced auditory neuronal and hair cell toxicity. *Laryngoscope.*2001;111(7):1147-55.
35. Tepel M. N-Acetylcysteine in the prevention of ototoxicity. *Kidney Int.*2007;72(3):231-2.
36. Schacht J. Antioxidant therapy attenuates aminoglycoside-induced hearing loss. *Ann N Y Acad Sci.*1999;884:125-30.
37. Husain K, Whitworth C, Somani SM, Rybak L P. Partial protection by lipoic acid against carboplatin-induced ototoxicity in rats. *Biomed Environ Sci.*2005;18(3):198-206.
38. Raponi G, Alpini D, Volonte S, Capobianco S, Cesarani A. The role of free radicals and plas-matic antioxidant in Meniere's syndrome. *Int Tinnitus J.*2003;9(2):104-8.
39. Savastano M, Brescia G, Marioni G. Antioxidant therapy in idiopathic tinnitus: preliminary outcomes. *Arch Med Res.* 2007;38(4):456-9.
40. Chen Y, Huang WG, Zha DJ, et al. Aspirin attenuates gentamicin ototoxicity: from the laboratory to the clinic. *Hear Res.*2007;226(1-2):178-82.
41. Shi X, Nuttall AL. Expression of adhesion molecular proteins in the cochlear lateral wall of normal and PARP-1 mutant mice. *Hear Res.* 2007;224(1-2):1-14.
42. Wang X, Truong T, Billings PB, Harris J P, Keithley EM. Blockage of immune-mediated inner ear damage by etanercept. *Otol Neurotol.* 2003;24(1):52-7
43. Derebery MJ. Allergic and immunologic aspects of Meniere's disease. *Otolaryngol Head Neck Surg.* 1996;114(3):360-5.
44. Prasad KN, Cole WC, Prasad KC. Risk factors for Alzheimer's disease: role of multiple antioxidants, non-steroidal anti-inflammatory and cholinergic agents alone or in combination in prevention and treatment. *J Am Coll Nutr.* 2002;21(6):506-22.
45. Prasad KN. Antioxidants in cancer care: when and how to use them as an adjunct to standard and experimental therapies. *Expert Rev Anticancer Ther.* 2003;3(6):903-15. 

Dr. Sanjay Agrawal,
 Leading Pharmaceutical Consultant
 and Editor-in Chief of *IJM Today*
 Post Graduation Diploma in
 Naturopathy and Yoga,
 6/146, Malviya Nagar,
 Jaipur -302017 ,
 Rajasthan